

2014年 英文誌掲載論文

APOE polymorphism and diabetic nephropathy.	Shin-ichi Araki	Clin Exp Nephrol.
High-density lipoprotein particle concentration and subclinical atherosclerosis of the carotid arteries in Japanese men.	Zaid M, Fujiyoshi A, Miura K, Abbott RD, Okamura T, Takashima N, Torii S, Saito Y, Hisamatsu T, Miyagawa N, Ohkubo T, Kadota A, Sekikawa A, Maegawa H, Nakamura Y, Mitsunami K, Ueshima H; For the SESSA Research group.	Atherosclerosis
Use of MRI signal intensity of extraocular muscles to evaluate methylprednisolone pulse therapy in thyroid-associated ophthalmopathy.	Higashiyama T, Nishida Y, Morino K, Ugi S, Nishio Y, Maegawa H, Ohji M.	Jpn J Ophthalmol.
Oral glucose-stimulated serum C-peptide predicts successful switching from insulin therapy to liraglutide monotherapy in Japanese patients with type 2 diabetes and renal impairment.	Araki H, Tanaka Y, Yoshida S, Morita Y, Kume S, Isshiki K, Araki S, Uzu T, Kashiwagi A, Maegawa H.	J Diabetes Investig.
Dapagliflozin as Monotherapy or Combination Therapy in Japanese Patients with Type 2 Diabetes: an Open-Label Study.	Kaku K, Maegawa H, Tanizawa Y, Kiyosue A, Ide Y, Tokudome T, Hoshino Y, Yang J, Langkilde AM.	Diabetes Ther.
Comparison of HOMA-IR, HOMA-β% and disposition index between US white men and Japanese men in Japan: the ERA JUMP study.	Ahuja V, Kadowaki T, Evans RW, Kadota A, Okamura T, El Khoudary SR, Fujiyoshi A, Barinas-Mitchell EJ, Hisamatsu T, Vishnu A, Miura K, Maegawa H, El-Saed A, Kashiwagi A, Kuller LH, Ueshima H, Sekikawa A; for the ERA JUMP Study Group.	Diabetologia.
A mutation of COX6A1 causes a recessive axonal or mixed form of Charcot-Marie-Tooth disease.	Tamiya G, Makino S, Hayashi M, Abe A, Numakura C, Ueki M, Tanaka A, Ito C, Toshimori K, Ogawa N, Terashima T, Maegawa H, Yanagisawa D, Tooyama I, Tada M, Onodera O, Hayasaka K.	Am J Hum Genet.
Role of nutrient-sensing signals in the pathogenesis of diabetic nephropathy.	Kume S, Koya D, Uzu T, Maegawa H	Biomed Res Int.
A high-fiber, low-fat diet improves periodontal disease markers in high-risk subjects: a pilot study.	Kondo K, Ishikado A, Morino K, Nishio Y, Ugi S, Kajiwara S, Kurihara M, Iwakawa H, Nakao K, Uesaki S, Shigeta Y, Imanaka H, Yoshizaki T, Sekine O, Makino T, Maegawa H, King GL, Kashiwagi A.	Nutr Res.
Predictive properties of plasma amino acid profile for cardiovascular disease in patients with type 2 diabetes.	Kume S, Araki S, Ono N, Shinhara A, Muramatsu T, Araki H, Isshiki K, Nakamura K, Miyano H, Koya D, Haneda M, Ugi S, Kawai H, Kashiwagi A, Uzu T, Maegawa H.	PLoS One.
Stem cell factor-activated bone marrow ameliorates amyotrophic lateral sclerosis by promoting protective microglial migration.	Terashima T, Kojima H, Urabe H, Yamakawa I, Ogawa N, Kawai H, Chan L, Maegawa H.	J Neurosci Res.
A fish-based diet intervention improves endothelial function in postmenopausal women with type 2 diabetes mellitus: a randomized crossover trial.	Kondo K, Morino K, Nishio Y, Kondo M, Nakao K, Nakagawa F, Ishikado A, Sekine O, Yoshizaki T, Kashiwagi A, Ugi S, Maegawa H.	Metabolism
Fatty acids are novel nutrient factors to regulate mTORC1 lysosomal localization and apoptosis in podocytes.	Yasuda M, Tanaka Y, Kume S, Morita Y, Chin-Kanasaki M, Araki H, Isshiki K, Araki S, Koya D, Haneda M, Kashiwagi A, Maegawa H, Uzu T.	Biochim Biophys Acta.
Long chain n-3 polyunsaturated fatty acids and incidence rate of coronary artery calcification in Japanese men in Japan and white men in the USA: population based prospective cohort study.	Sekikawa A, Miura K, Lee S, Fujiyoshi A, Edmundowicz D, Kadowaki T, Evans RW, Kadowaki S, Sutton-Tyrrell K, Okamura T, Bertolet M, Masaki KH, Nakamura Y, Barinas-Mitchell EJ, Willcox BJ, Kadota A, Seto TB, Maegawa H, Kuller LH, Ueshima H; ERA JUMP Study Group.	Heart.
CCDC3 is specifically upregulated in omental adipose tissue in subjects with abdominal obesity.	Ugi S, Maeda S, Kawamura Y, Kobayashi MA, Imamura M, Yoshizaki T, Morino K, Sekine O, Yamamoto H, Tani T, Rokushima M, Kashiwagi A, Maegawa H	Obesity (Silver Spring)
Safety and efficacy of skin patches containing loxoprofen sodium in diabetic patients with overt nephropathy.	Araki H, Kuwagata S, Soumura M, Yamahara K, Morita Y, Kume S, Isshiki K, Araki SI, Kashiwagi A, Maegawa H, Uzu T.	Clin Exp Nephrol.