

2020年 英文誌掲載論文

A Low-Carbohydrate Diet Improves Glucose Metabolism in Lean Insulinopenic Akita Mice Along With Sodium-Glucose Cotransporter 2 Inhibitor	Fujita Y, Atageldiyeva KK, Takeda Y, Yanagimachi T, Makino Y, Haneda M.	Front Endocrinol (Lausanne).
Safety and effectiveness of ipragliflozin in Japanese patients with type 2 diabetes mellitus and impaired renal function: subgroup analysis of a 3-year post-marketing surveillance study (STELLA-LONG TERM)	Tobe K, Maegawa H, Nakamura I, Uno S.	Diabetol Int.
Impact of obesity on underreporting of energy intake in type 2 diabetic patients: Clinical Evaluation of Energy Requirements in Patients with Diabetes Mellitus (CLEVER-DM) study	Miyazawa I, Morino K, Fuse K, Kondo K, Ohi A, Nishida K, Kurihara M, Yasuhara S, Nakanishi N, Nishida Y, Nakae S, Yamada Y, Tanaka S, Ebine N, Sasaki S, Katsukawa F, Sasaki M, Ugi S, Maegawa H; CLEVER-DM Study Research Group.	Clin Nutr ESPEN.
SGLT2 Inhibition Mediates Protection from Diabetic Kidney Disease by Promoting Ketone Body-Induced mTORC1 Inhibition	Tomita I, Kume S, Sugahara S, Osawa N, Yamahara K, Yasuda-Yamahara M, Takeda N, Chin-Kanasaki M, Kaneko T, Mayoux E, Mark M, Yanagita M, Ogita H, Araki SI, Maegawa H.	Cell Metab.
Family Support for Medical Nutritional Therapy and Dietary Intake among Japanese with Type 2 Diabetes (JDDM 56)	Horikawa C, Hatta M, Morikawa SY, Takeda Y, Takeuchi M, Fujihara K, Kato N, Yokoyama H, Kurihara Y, Iwasaki K, Tanaka S, Maegawa H, Sone H.	Nutrients.
Contrast medium-induced severe thrombocytopenia in patient on maintenance hemodialysis: a case report and literature review.	Takeda N, Araki SI, Chin-Kanasaki M, Osawa N, Sawai K, Yamahara K, Yasuda-Yamahara M, Kume S, Fujita Y, Maegawa H.	CEN Case Rep.
A Real-World Observational Study Evaluating the Probability of Glycemic Control with Basal Insulin or Glucagon-Like Peptide-1 Receptor Agonist in Japanese Patients with Type 2 Diabetes	Baxter M, Morimoto Y, Tamiwa M, Hattori M, Peng XV, Lubwama R, Maegawa H.	Diabetes Ther.
The Prognosis of Patients With Type 2 Diabetes and Nonalbuminuric Diabetic Kidney Disease Is Not Always Poor: Implication of the Effects of Coexisting Macrovascular Complications (JDDM 54).	Yokoyama H, Araki SI, Kawai K, Yamazaki K, Shirabe SI, Sugimoto H, Minami M, Miyazawa I, Maegawa H; JDDM Study Group.	Diabetes Care.
Protective role of podocyte autophagy against glomerular endothelial dysfunction in diabetes.	Yoshibayashi M, Kume S, Yasuda-Yamahara M, Yamahara K, Takeda N, Osawa N, Chin-Kanasaki M, Nakae Y, Yokoi H, Mukoyama M, Asanuma K, Maegawa H, Araki SI.	Biochem Biophys Res Commun.
Validity of the Use of a Triaxial Accelerometer and a Physical Activity Questionnaire for Estimating Total Energy Expenditure and Physical Activity Level among Elderly Patients with Type 2 Diabetes Mellitus: CLEVER-DM Study.	Nishida Y, Tanaka S, Nakae S, Yamada Y, Morino K, Kondo K, Nishida K, Ohi A, Kurihara M, Sasaki M, Ugi S, Maegawa H, Ebine N, Sasaki S, Katsukawa F.	Ann Nutr Metab
Liver fat accumulation assessed by computed tomography is an independent risk factor for diabetes mellitus in a population-based study: SESSA (Shiga Epidemiological Study of Subclinical Atherosclerosis).	Fuse K, Kadota A, Kondo K, Morino K, Fujiyoshi A, Hisamatsu T, Kadowaki S, Miyazawa I, Ugi S, Maegawa H, Miura K, Ueshima H; SESSA Research Group.	Diabetes Res Clin Pract.
Relationship between carbohydrate and dietary fibre intake and the risk of cardiovascular disease mortality in Japanese: 24-year follow-up of NIPPON DATA80	Miyazawa I, Miura K, Miyagawa N, Kondo K, Kadota A, Okuda N, Fujiyoshi A, Chihara I, Nakamura Y, Hozawa A, Nakamura Y, Kita Y, Yoshita K, Okamura T, Okayama A, Ueshima H; NIPPON DATA80 Research group.	Eur J Clin Nutr.
Combined effect of sodium-glucose cotransporter 2 and dipeptidyl peptidase-4 inhibitors for diabetic kidney disease.	Araki SI.	J Diabetes Investig.